PubMed	
Format: Abstract	Full text links
Fur J Epidemiol, 2016 May:31(5):491-9, doi: 10.1007/s1065	SpringerLink PMC Full text

Low serum vitamin D is associated with axial length and risk of myopia in young children.

<u>Tideman JW</u>^{1,2}, <u>Polling JR</u>^{1,3}, <u>Voortman T</u>², <u>Jaddoe VW</u>^{2,4}, <u>Uitterlinden AG</u>^{2,5}, <u>Hofman A</u>², <u>Vingerling JR</u>¹, <u>Franco OH</u>², <u>Klaver CC</u>^{6,7}.

Author information


Abstract

The aim of the study was to investigate the relationship between serum 25(OH)D levels and axial length (AL) and myopia in 6-year-old children. A total of 2666 children aged 6 years participating in the birth-cohort study Generation R underwent a stepwise eye examination. First, presenting visual acuity (VA) and AL were performed. Second, automated cycloplegic refraction was measured if LogMAR VA > 0.1. Serum 25hydroxyvitamin D [25(OH)D] was determined from blood using liquid chromatography/tandem mass spectrometry. Vitamin D related SNPs were determined with a SNP array; outdoor exposure was assessed by questionnaire. The relationships between 25(OH)D and AL or myopia were investigated using linear and logistic regression analysis. Average 25(OH)D concentration was 68.8 nmol/L (SD ± 27.5; range 4-211); average AL 22.35 mm (SD ± 0.7; range 19.2-25.3); and prevalence of myopia 2.3 % (n = 62). After adjustment for covariates, 25(OH)D concentration (per 25 nmol/L) was inversely associated with AL (β -0.043; P < 0.01), and after additional adjusting for time spent outdoors (β -0.038; P < 0.01). Associations were not different between European and non-European children (β -0.037 and β -0.039 respectively). Risk of myopia (per 25 nmol/L) was OR 0.65 (95 % CI 0.46-0.92). None of the 25(OH)D related SNPs showed an association with AL or myopia. Lower 25(OH)D concentration in serum was associated with longer AL and a higher risk of myopia in these young children. This effect appeared independent of outdoor exposure and may suggest a more direct role for 25(OH)D in myopia pathogenesis.

KEYWORDS: Axial length; Children; Myopia; Vitamin D

PMID: 26955828 PMCID: PMC4901111 DOI: 10.1007/s10654-016-0128-8

[Indexed for MEDLINE] Free PMC Article



MeSH terms, Substances

LinkOut - more resources